

# Demographic Variations in Maxillary Pathologic Lesions - An Institutional Study

**Shivangi Gaur**

*Department of Oral & Maxillofacial Surgery  
Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical And Technical Science,  
Saveetha University, Chennai- 600077, India.  
Email address- 151804004.sdc@saveetha.com*

**Madhulaxmi M**

Professor

*Department of Oral & Maxillofacial Surgery  
Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical And Technical Science,  
Saveetha University, Chennai- 600077, India.  
Email address- madhulaxmi@saveetha.com*

**Hemavathy O R**

Associate Professor

*Department of Oral & Maxillofacial Surgery  
Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical And Technical Science,  
Saveetha University, Chennai- 600077, India.  
Email address-hemavathy.sdc@saveetha.com*

**Corresponding Author**

**Madhulaxmi M**

Professor

*Department of Oral & Maxillofacial Surgery  
Saveetha Dental College and Hospitals,  
Saveetha Institute of Medical And Technical Science,  
Saveetha University, Chennai- 600077, India  
Email address- madhulaxmi@saveetha.com*

**Article Info**

**Volume 83**

**Page Number: 2409 - 2424**

**Publication Issue:**

**July-August 2020**

**Abstract:**

A variety of benign and malignant lesions involve the maxilla and maxillary sinus. The presenting features, symptomatology and advanced diagnostic imaging techniques help in achieving a differential diagnosis however histopathological examination remains the mainstay of definitive diagnosis. This study aims to identify the frequency of isolated maxillary pathologic lesions (benign and malignant) and its gender and age distribution. After clearance from the institutional ethical committee, a retrospective study was performed for a period of ten months from June 2019 to March 2020 in our institution. 90 cases of pathology were reviewed out of which 29 were included in the study based on the inclusion criteria. 68.97% males and 31.03% females reported with pathologic lesions of the maxilla alone in our center. Malignant maxillary lesions show the highest frequency of occurrence in 17.24% in the 41-60 age group (p value-0.233). Treatment modalities used for various lesions diagnosed in the maxilla showed the highest degree of association with surgical excision being the most commonly performed for malignant lesions and enucleation for cysts and benign tumors (p value- 0.000). Maxillary benign pathological lesions have a tendency to mimic malignant lesions. Hence it is important to document and identify lesions most commonly encountered in a particular population to provide for adequate and appropriate treatment care needs.

**Keywords:** Rural Maxilla; Pathology; Malignant; Benign; Cysts; Tumors

**Article History**

**Article Received:06 June 2020**

**Revised: 29 June 2020**

**Accepted: 14 July 2020**

**Publication: 25 July 2020**

## INTRODUCTION

Maxilla has a complex structure and is the most important bone of the midface. Due to its central location it provides functional support to the viscerocranium. It also plays a fundamental role in facial architecture, separating nasal and oral cavities, forms the upper jaw and contains the maxillary sinus (Dalgorf and Higgins, 2008; Fehrenbach and Herring, 2013; Kühnel and Reichert, 2015). Each maxillary body contains an air filled hollow cavity called the maxillary sinus which is the seat of resonance of voice and provides pneumatization of maxilla thereby reducing the weight of the skull (Cunningham, 1818; Larrabee, Makielski and Henderson, 2004; Lev and Artzi, 2020; Morris and Tadi, 2020).

The lesions encountered in the maxilla or mandible include cysts of odontogenic and non-odontogenic origin, tumors of odontogenic origin, giant cell lesions, salivary gland diseases and tumors, vascular anomalies, osteomyelitis, osteoradionecrosis, secondarily infected cystic lesions, benign and

malignant neoplasms. (Regezi, Sciubba and Jordan, 2012; Carton, 2017; Thompson and Bishop, 2017; Alramadhan *et al.*, 2020; Banasser *et al.*, 2020). Epidemiological data for pathological lesions affecting mandible or maxilla is helpful in determination of demographic characteristics of oral lesions.

There are numerous pathological lesions affecting the maxilla and maxillary sinus, treatment of which leaves a midfacial defect which needs to be addressed while forming a treatment plan. Predominantly the pathological lesions found to be reported in the literature majorly in maxilla include - Adenomatoid odontogenic tumor (Adisa *et al.*, 2016; Dhupar, Akkara and Khandelwal, 2016; Katiyar *et al.*, 2019; Mohanty *et al.*, 2019), Ossifying Fibroma (Thakur *et al.*, 2011; George *et al.*, 2017; Muthu, 2017; Seo *et al.*, 2019; Almeida Júnior *et al.*, 2020; Macedo *et al.*, 2020), Odontoma (Christensen, 1956; Wanjari, Wanjari and Bhowate, 2011; Murphy *et al.*, 2014; Lee *et al.*, 2016; Kalra *et al.*, 2018; de Oliveira *et al.*, 2019; Zhuoying and Fengguo, 2019), Fibrous dysplasia (Ward, Alley and Owen, 1969;

'Monostotic Type of Fibrous Dysplasia of Maxilla', 2016; Kochanowski *et al.*, 2018; Shanmugam *et al.*, 2018; Su *et al.*, 2019), Paget's disease (Mooney *et al.*, 2003; Karakida *et al.*, 2010), ghost cell odontogenic carcinoma (de Arruda *et al.*, 2018), surgical ciliated cyst (Leung, Wong and Cheung, 2012; Chee *et al.*, 2014; Nogueira *et al.*, 2014), osteomyelitis (Himalstein, 1967; Meethal, 2016; Pattnaik *et al.*, 2017; Antunes *et al.*, 2018; Arani, Shareef and Khanam, 2019; Sadaksharam and Murugesan, 2019), osteonecrosis (Seehra, Sloan and Oliver, 2009; Kolar, Nair and Kumar, 2015).

The sparsity of epidemiological data on pathological lesions predominantly affecting the maxilla was the reason for formulating this study in a south indian sample population. Till date the institutional team has conducted several clinical trials (Jesudasan, Wahab and Sekhar, 2015; Christabel *et al.*, 2016; Kumar, 2017b; Packiri, Gurunathan and Selvarasu, 2017; Patil *et al.*, 2017; Rao and Santhosh Kumar, 2018; Abhinav *et al.*, 2019; Jain *et al.*, 2019; Sweta, Abhinav and Ramesh, 2019), genetic studies (Marimuthu *et al.*, 2018), various surveys (Kumar and Sneha, 2016; Patturaja and Pradeep, 2016; Kumar, 2017a; Kumar and Rahman, 2017) and literature update (Kumar, 2017c) over various topics of surgical interest. The aim of this study was to identify the frequency of isolated maxillary pathologic lesions (benign and malignant) and its gender and age distribution.

## MATERIALS AND METHODS

A retrospective study on prevalence of maxillary pathological lesions treated under general anesthesia was done in our institution over a period of ten months from June 2019 to March 2020. Institutional ethical clearance was obtained for data retrieval and usage as required for the study (SDC/SIHEC/2020/DIASDATA/0619-0320).

A total of 485 patients had undergone surgical intervention under general anesthesia during the

study period. Among them ninety patients were treated for benign or malignant pathologies. The other treatments including those for trauma, congenital deformities, cleft lip/palate, orthognathic corrective surgeries, TMJ surgeries, preprosthetic surgeries and disimpactions were excluded. Among the ninety case sheets of patients undergoing treatment for pathological lesions, sixty one pathologies were in other anatomical sites not involving maxilla. Case sheets of all relevant patients with pathologies involving maxilla alone exclusively were identified and reviewed for inclusion in the study. After taking into consideration the inclusion and exclusion criteria, it was found that a total of 29 patients met our inclusion criteria.


All patients from 0 to 80 years old irrespective of their medical conditions who underwent primary surgical management of histopathologically proven maxillary pathological lesions were included as study subjects. The lesions included were benign or malignant tumors of maxilla, minor salivary gland pathology involving maxilla, developmental or inflammatory cysts in the maxilla which had complete preoperative, intraoperative and postoperative data available. Exclusion criteria were all pathological lesions involving mandible, buccal mucosa, tongue, major salivary gland pathologies, congenital deformities and incomplete data.

Data collection was performed to include demographic details along with the histopathological diagnosis and treatment done. Parameters evaluated were age, gender, diagnosis, treatment done dependent variable and maxillary pathological lesions with the independent variable. Data was analyzed using SPSS version 20 statistical software. The collected data was analysed using SPSS software version 20. Chi square test was used and p - value of <0.05 was taken as statistically significant.

## RESULTS AND DISCUSSION

Upon data analysis, there were a total of 29 patients out of 90 identified (32%) as isolated maxillary


pathological lesions treated under general anaesthesia. Out of which 68.97% (n=20) were males and 31.03% (n=9) were females (Graph 1)



Graph 1 : Bar graph showing the distribution of maxillary pathological lesions amongst the genders. 68.97% males (blue) and 31.03% females (brown) reported with pathologic lesions of the maxilla in our center

which is the general trend worldwide (Ghazawi *et al.*, 2020; Kammer, Mello and Rivero, 2020). Amongst males the lesions involving the maxilla reported with the highest frequency of 20.69% (n=6) were malignant tumors, followed closely by inflammatory lesions including osteomyelitis 17.24% (n=5), 13.79% (n=4) of developmental cysts, 10.34% (n=3) of inflammatory cysts and 6.90%

(n=2) benign tumors. Females present with the highest frequency of inflammatory cysts of 13.79% (n=4) and equal frequencies of 6.90% (n=2) benign and malignant lesions, with least reported lesion being developmental cysts with a frequency of 3.45% (n=1) (Graph 2).




Graph 2 : Bar graph illustrating the relationship of gender with the lesions diagnosed in the maxilla where X axis represents gender, Y axis represents total number of patients treated for maxillary pathologic lesions.

Males report with the highest frequency of 20.69% malignant tumors (violet), followed by 17.24% inflammatory lesions (yellow). Females present with the highest frequency of 13.79% inflammatory cysts (green). These associations are however not significant statistically ( p value-0.233 on Chi Square test)

Males are generally more commonly affected by malignancies such as squamous cell carcinoma(Serindere *et al.*, 2019; Ghazawi *et al.*, 2020), developmental and inflammatory cysts (Tamiolakis *et al.*, 2019; Kammer, Mello and Rivero, 2020) which was consistent with our study. However, in the population studied by (Kilinc *et al.*, 2017; Silva *et al.*, 2018; Farias *et al.*, 2019) females were more commonly affected by developmental cysts and tumors, which was not the same in our

study population which could be due to certain reasons like, asymptomatic lesions going unnoticed and undetected for several years in the absence of imaging by the clinician or in the absence of any clinical signs - like expansion of bone, mobility of tooth, paresthesia etc to suggestive of a pathology (Dong *et al.*, 2020; Goto *et al.*, 2020; Li *et al.*, 2020; MacDonald, Martin and Nguyen, 2020; Maruyama *et al.*, 2020; Shimmura *et al.*, 2020; Xia *et al.*, 2020).




Graph 3 : Bar graph showing the distribution of maxillary pathological lesions amongst the different age groups, where X axis represents the age and Y axis represents the number of patients. 21-40 age group individuals (teal) reported with the highest frequency of maxillary pathological lesions 37.93% , followed by 41-60 age group individuals (grey) with 34.48% lesions, >60 age group individuals with 17.24% lesions. 10.34 % lesions reported were reported in the <20 age group (light blue)

Graph 3 shows the distribution of maxillary pathological lesions amongst the different age groups, divided as below 20 years old, 21-40 years old, 41-60 years old and above 60 years old for the purpose of this study. 21-40 age group individuals reported with the highest frequency of maxillary pathological lesions 37.93% (n=11); followed by 41-60 age group individuals with 34.48% (n=10) lesions, below 60 age group individuals with 17.24% (n=5) maxillary pathologic lesions and 10.34% (n=3) pathologic maxillary lesions reported in the

below 20 age group. According to literature, most commonly pathologic lesions are presented in 4 to 6th decade of life(Tamirakis *et al.*, 2019). However, in our study there was an almost equal frequency of maxillary pathological lesion seen from the second to fourth decade .

We also studied the relationship of the most common lesions of the maxilla being diagnosed in particular age groups (Graph 4).




Graph 4 : Bar graph illustrating relationship of age with the lesions diagnosed in the maxilla where X axis represents age group, Y axis represents total number of patients treated for maxillary pathologic lesions. Malignant lesions (violet) show the highest frequency of 17.24% in 41-60 and 10.34% in >60 age groups. Inflammatory cysts (green) occur 10.34% in the 21-40 age group. These associations are however not significant statistically ( p value-0.211 on Chi Square test)


In the 20 years age group, all three lesions that is developmental cysts, inflammatory lesions and benign tumors of maxilla - were diagnosed with an equal frequency of 3.45% (n=1). 21-40 age group shows the highest frequency of being diagnosed with inflammatory lesions of the maxilla with a frequency of 13.79% (n=4), followed by inflammatory cysts with a frequency of 10.34% (n=3), with an equal frequency of 6.90% (n=2) being diagnosed with developmental cysts and benign tumors. However, the bulk of maxillary pathologic lesions, that is malignant lesions are found in the 41-60 year olds with a frequency of 17.24% (n=5) in our study, followed by equal frequency of 6.90% (n=2) for diagnosis of developmental or inflammatory cysts

and least frequency of benign tumor of 3.45% (n=1) in this age group. According to previously published literature patients presenting with developmental cysts were older than 20 years of age, which is consistent with our results (Sriram and Shetty, 2008; Luo and Li, 2009; Rajendran and Sivapathasundharam, 2014).

We studied the distribution of maxillary pathological lesions amongst the sample population. 27.59% (n=8) were malignant tumors , 24.14% (n=7) inflammatory cysts , 17.24% (n=5) developmental cysts, 17.24% (n=5) inflammatory lesions and 13.79% (n=4) benign tumors of the maxilla were diagnosed in our study (Graph 5).




Graph 5 : Bar graph showing the distribution of maxillary pathological lesions amongst the sample population, where X axis represents the type of pathology and Y axis represents the number of patients. Lesions are distributed as 27.59% malignant tumors (violet), 24.14% inflammatory cysts (green), 17.24% developmental cysts (cobalt blue) and inflammatory lesions (yellow) respectively and 13.79% benign tumors (beige)



Graph 6 : Bar graph showing the distribution of treatment done for maxillary pathological lesions amongst the sample population, where X axis represents the varied treatment and Y axis the number of patients. Enucleation (black) is the most commonly used treatment modality with a frequency of 62.07%, followed by surgical enucleation (light grey) with a frequency of 31.03% and equal frequency of 3.45% marsupialization (pink) and sequestrectomy (green).

Graph 6 shows the distribution of treatment done where enucleation is the most commonly used treatment modality with a frequency of 62.07% (n=18), followed by surgical enucleation 31.03% (n=9) and equal frequency of 3.45% (n=1) marsupialization and sequestrectomy. The association of diagnosis and treatment are significant statistically with a p value-0.000 on Chi Square test. Malignant lesions were treated by using surgical

excision most commonly in all 27.59% (n=8) cases. Inflammatory cysts were treated by enucleation in all 24.14% (n=7) cases. Benign lesions were treated using both marsupialisation and surgical excision with equal frequency of 3.45% (n=1). Inflammatory lesions were treated using sequestrectomy in 3.45%(n=1) and enucleation in 13.79% (n=4) cases (Graph 7).



Graph 7 : Bar graph illustrating the spectrum of treatment performed for various lesions diagnosed in the maxilla, where X axis represents treatment, Y axis represents total number of patients treated for maxillary pathologic lesions. Surgical excision is the most commonly performed treatment 27.59% for malignant lesions (violet). Inflammatory cysts (green) are treated by enucleation in all 24.14% cases. The association between the type of pathology and the treatment done was highly significant statistically with p value =  $0.000 < 0.05$  on Chi Square test.

In our knowledge this study is the first of its kind to describe oral pathological lesions with site/location as the primary variable of interest followed by frequency of these lesions reported in the maxillary arch alone. Due to the complex anatomy and function of maxilla, it is important to define the commonly encountered lesions in a given population over a certain period of time.

The study endeavoured to collect epidemiological data from a sample south Indian population to study the prevalence or frequency of involvement of maxilla in oral pathological lesions and as consistent with the literature, the maxilla is less frequently

involved, but lesions become more aggressive and require more aggressive treatment course due to close affinity with the surrounding structures such as nasal cavity, antrum of highmore, cavernous sinus, etc. Also all lesions associated with extracted teeth like radicular cyst, pyogenic granuloma etc may not be reported to the histopathologist, leading to some deficiency in number of oral pathological lesions in the maxilla. We believe that epidemiological analysis can be continued for a longer duration of time and using a wider community to acquire certain definitive perspectives on the lesions involving the maxilla. Maxillary lesions, although infrequent and rarer than mandibular lesions, require aggressive

treatment which can be debilitating due to the close approximation with vital anatomic structures thereby compelling prolonged follow up.

## CONCLUSION

This study concludes that more than one quarter of the pathologic lesions identified in the oral and maxillofacial region were isolated maxillary pathologies with a gender predilection to males. Benign pathologies were more common in the second to third decade and malignant pathologies more common in the fourth to sixth decade of life. However, malignant pathologies encountered in maxilla were comparable to or more than benign and inflammatory pathologies. This emphasises the need for early diagnosis and treatment protocols without undue delay in maxillary pathologies. Due to the occult nature of lesions of the maxilla, it is difficult to detect these at an early stage. Hence, identification of most commonly encountered lesions affecting the maxilla in a particular geographic distribution will aid the clinician to formulate diagnosis with ease and predictability and choose appropriate treatment course depending upon the needs of the individual.

## REFERENCES

1. Abhinav, R. P. *et al.* (2019) 'The Patterns and Etiology of Maxillofacial Trauma in South India', *Annals of maxillofacial surgery*, 9(1), pp. 114–117.
2. Adisa, A. O. *et al.* (2016) 'A retrospective review of 61 cases of adenomatoid odontogenic tumour seen in five tertiary health facilities in Nigeria', *The Pan African medical journal*, 24, p. 102.
3. Almeida Júnior, V. R. de *et al.* (2020) 'Simultaneous presentation of juvenile ossifying fibroma in the maxilla and mandible: a case report', *International journal of surgery case reports*, 71, pp. 285–289.
4. Alramadhan, S. A. *et al.* (2020) 'Retrospective Study of Buccal Mucosal Salivary Neoplasms', *Head and neck pathology*. doi: 10.1007/s12105-020-01190-8.
5. Antunes, L. de O. *et al.* (2018) 'Osteomyelitis of the maxilla caused by sp', *Radiologia brasileira*, 51(5), pp. 340–341.
6. Arani, R., Shareef, S. N. H. A. and Khanam, H. M. K. (2019) 'Mucormycotic Osteomyelitis Involving the Maxilla: A Rare Case Report and Review of the Literature', *Case reports in infectious diseases*, 2019, p. 8459296.
7. de Arruda, J. A. A. *et al.* (2018) 'Calcifying odontogenic cyst, dentinogenic ghost cell tumor, and ghost cell odontogenic carcinoma: A systematic review', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 47(8), pp. 721–730.
8. Banasser, A. M. *et al.* (2020) 'Odontogenic Myxoma: A 23-Year Retrospective Series of 38 Cases', *Head and neck pathology*. doi: 10.1007/s12105-020-01191-7.
9. Carton, J. (2017) 'Head and neck pathology', *Oxford Medicine Online*. doi: 10.1093/med/9780198759584.003.0006.
10. Chee, C. A. K. M. *et al.* (2014) 'Surgical ciliated cyst of the maxilla following maxillary osteotomy: a case report', *Oral Surgery*, pp. 39–41. doi: 10.1111/ors.12041.
11. Christabel, A. *et al.* (2016) 'Comparison of pterygomaxillary dysjunction with tuberosity separation in isolated Le Fort I osteotomies:

- a prospective, multi-centre, triple-blind, randomized controlled trial', *International journal of oral and maxillofacial surgery*, 45(2), pp. 180–185.
12. Christensen, R. W. (1956) 'Complex composite odontoma involving the maxilla and maxillary sinus', *Oral Surgery, Oral Medicine, Oral Pathology*, pp. 1156–1164. doi: 10.1016/0030-4220(56)90185-2.
  13. Cunningham, D. J. (1818) *Cunningham's textbook of anatomy*.
  14. Dalgorf, D. and Higgins, K. (2008) 'Reconstruction of the midface and maxilla', *Current opinion in otolaryngology & head and neck surgery*, 16(4), pp. 303–311.
  15. Dhupar, V., Akkara, F. and Khandelwal, P. (2016) 'An unusually large aggressive adenomatoid odontogenic tumor of maxilla involving the third molar: A clinical case report', *European journal of dentistry*, 10(2), pp. 277–280.
  16. Dong, D. *et al.* (2020) 'Fibro-osseous Lesions of Paranasal Sinus and Craniofacial Region: A Retrospective Study of 282 Cases', *The Laryngoscope*. doi: 10.1002/lary.28529.
  17. Farias, J. G. *et al.* (2019) 'Epidemiological study of intraosseous lesions of the stomatognathic or maxillomandibular complex diagnosed by a Reference Centre in Brazil from 2006-2017', *The British journal of oral & maxillofacial surgery*, 57(7), pp. 632–637.
  18. Fehrenbach, M. J. and Herring, S. W. (2013) *Illustrated Anatomy of the Head and Neck - E-Book*. Elsevier Health Sciences.
  19. George, S. *et al.* (2017) 'Giant Peripheral Ossifying Fibroma of Maxilla', *Indian Journal of Dental Education*, pp. 74–77. doi: 10.21088/ijde.0974.6099.10217.15.
  20. Ghazawi, F. M. *et al.* (2020) 'Epidemiology and Patient Distribution of Oral Cavity and Oropharyngeal SCC in Canada', *Journal of cutaneous medicine and surgery*, p. 1203475420915448.
  21. Goto, M. *et al.* (2020) 'A rare case of odontogenic keratocyst extending into the sphenoid bone from the maxilla', *International journal of surgery case reports*, 71, pp. 132–138.
  22. Himalstein, M. R. (1967) 'OSTEOMYELITIS OF THE MAXILLA DUE TO AN ANAEROBIC ORGANISM', *The Laryngoscope*, pp. 559–561. doi: 10.1288/00005537-196704000-00011.
  23. Jain, S. V. *et al.* (2019) 'Evaluation of Three-Dimensional Changes in Pharyngeal Airway Following Isolated Lefort One Osteotomy for the Correction of Vertical Maxillary Excess: A Prospective Study', *Journal of Maxillofacial and Oral Surgery*, pp. 139–146. doi: 10.1007/s12663-018-1113-4.
  24. Jesudasan, J. S., Wahab, P. U. A. and Sekhar, M. R. M. (2015) 'Effectiveness of 0.2% chlorhexidine gel and a eugenol-based paste on postoperative alveolar osteitis in patients having third molars extracted: a randomised controlled clinical trial', *The British journal of oral & maxillofacial surgery*, 53(9), pp. 826–830.
  25. Kalra, A. *et al.* (2018) 'A Bag of Marbles: A Fascinating Compound Odontoma of the Maxilla', *Pediatric dentistry*, 40(2), pp. 140–142.
  26. Kammer, P. V., Mello, F. W. and Rivero, E. R. C. (2020) 'Comparative analysis between developmental and inflammatory

- odontogenic cysts: retrospective study and literature review', *Oral and maxillofacial surgery*, 24(1), pp. 73–84.
27. Karakida, K. *et al.* (2010) 'Multiple giant cell tumors in maxilla and skull complicating Paget's disease of bone', *The Tokai journal of experimental and clinical medicine*, 35(3), pp. 112–117.
  28. Katiyar, A. *et al.* (2019) 'Trauma to Tumor: A Hunt of Adenomatoid Odontogenic Tumor-A Rare Case Report', *International journal of clinical pediatric dentistry*, 12(4), pp. 366–369.
  29. Kilinc, A. *et al.* (2017) 'Benign tumors and tumor-like lesions of the oral cavity and jaws: An analysis of 709 cases', *Nigerian journal of clinical practice*, 20(11), pp. 1448–1454.
  30. Kochanowski, N. E. *et al.* (2018) 'Radiographic Diagnosis of Fibrous Dysplasia in Maxilla', *Cureus*. doi: 10.7759/cureus.3127.
  31. Kolor, T., Nair, S. C. and Kumar, B. (2015) 'Osteonecrosis of maxilla secondary to bisphosphonate therapy: a case report', *Journal of maxillofacial and oral surgery*, 14(Suppl 1), pp. 52–56.
  32. Kühnel, T. S. and Reichert, T. E. (2015) 'Trauma of the midface', *GMS current topics in otorhinolaryngology, head and neck surgery*, 14, p. Doc06.
  33. Kumar, S. (2017a) 'KNOWLEDGE, ATTITUDE AND AWARENESS OF DENTAL UNDERGRADUATE STUDENTS REGARDING HIV/AIDS PATIENTS', *Asian Journal of Pharmaceutical and Clinical Research*, 10(5), p. 175.
  34. Kumar, S. (2017b) 'RELATIONSHIP BETWEEN DENTAL ANXIETY AND PAIN EXPERIENCE DURING DENTAL EXTRACTIONS', *Asian Journal of Pharmaceutical and Clinical Research*, p. 458. doi: 10.22159/ajpcr.2017.v10i3.16518.
  35. Kumar, S. (2017c) 'THE EMERGING ROLE OF BOTULINUM TOXIN IN THE TREATMENT OF OROFACIAL DISORDERS: LITERATURE UPDATE', *Asian Journal of Pharmaceutical and Clinical Research*, p. 21. doi: 10.22159/ajpcr.2017.v10i9.16914.
  36. Kumar, S. and Rahman, R. (2017) 'KNOWLEDGE, AWARENESS, AND PRACTICES REGARDING BIOMEDICAL WASTE MANAGEMENT AMONG UNDERGRADUATE DENTAL STUDENTS', *Asian Journal of Pharmaceutical and Clinical Research*, 10(8), p. 341.
  37. Kumar, S. and Sneha, S. (2016) 'KNOWLEDGE AND AWARENESS REGARDING ANTIBIOTIC PROPHYLAXIS FOR INFECTIVE ENDOCARDITIS AMONG UNDERGRADUATE DENTAL STUDENTS', *Asian Journal of Pharmaceutical and Clinical Research*, p. 154. doi: 10.22159/ajpcr.2016.v9s2.13405.
  38. Larrabee, W. F., Makielski, K. H. and Henderson, J. L. (2004) *Surgical Anatomy of the Face*. Lippincott Williams & Wilkins.
  39. Lee, H. M. *et al.* (2016) 'Compound Type Odontoma at Maxilla', *Archives of Craniofacial Surgery*, p. 96. doi: 10.7181/acfs.2016.17.2.96.
  40. Leung, Y. Y., Wong, W. Y. and Cheung, L. K. (2012) 'Surgical ciliated cysts may mimic radicular cysts or residual cysts of maxilla: 2421

- report of 3 cases', *Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons*, 70(4), pp. e264–9.
41. Lev, D. and Artzi, Z. (2020) 'The Anatomy of the Maxilla and the Mandible', *Bone Augmentation by Anatomical Region*, pp. 1–16. doi: 10.1002/9781119427926.ch1.
  42. Li, S. H. *et al.* (2020) 'Endoscope-assisted Surgery in the Treatment of Dentigerous Cyst Involving the Maxillary Sinus - Report of Two Cases', *The Chinese journal of dental research: the official journal of the Scientific Section of the Chinese Stomatological Association*, 23(1), pp. 71–76.
  43. Luo, H.-Y. and Li, T.-J. (2009) 'Odontogenic tumors: a study of 1309 cases in a Chinese population', *Oral oncology*, 45(8), pp. 706–711.
  44. MacDonald, D., Martin, M. and Nguyen, C. (2020) 'Malignant lesions in the anterior maxilla', *Clinical radiology*, 75(7), pp. 497–506.
  45. Macedo, D. de V. *et al.* (2020) 'Ossifying fibroma in the maxilla and orbital floor: report of an uncommon case', *Journal of the Korean Association of Oral and Maxillofacial Surgeons*, 46(3), pp. 204–207.
  46. Marimuthu, M. *et al.* (2018) 'Canonical Wnt pathway gene expression and their clinical correlation in oral squamous cell carcinoma', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(3), pp. 291–297.
  47. Maruyama, N. *et al.* (2020) 'Quadruple Multiple Primary Malignancies: Early Detection of Second Primary Malignancy by Esophagogastroduodenoscopy/Colonoscopy Is Crucial for Patients with Classic Kaposi's Sarcoma', *Diagnostics (Basel, Switzerland)*, 10(4). doi: 10.3390/diagnostics10040218.
  48. Meethal, A. C. (2016) 'Actinomycotic Osteomyelitis of the Maxilla – A Delusive Presentation', *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. doi: 10.7860/jcdr/2016/19171.8086.
  49. Mohanty, R. *et al.* (2019) 'A rare nonsyndromic case of adenomatoid odontogenic tumor associated with multiple impacted supernumerary teeth', *National journal of maxillofacial surgery*, 10(1), pp. 114–117.
  50. 'Monostotic Type of Fibrous Dysplasia of Maxilla' (2016) *Indian Journal of Dental Advancements*. doi: 10.5866/2016.8.10263.
  51. Mooney, W. W. W. *et al.* (2003) 'Recurrent giant cell tumour of the maxilla associated with both Paget's disease and primary hyperparathyroidism', *ANZ journal of surgery*, 73(10), pp. 863–864.
  52. Morris, A. L. and Tadi, P. (2020) 'Anatomy, Head and Neck, Teeth', in *StatPearls*. Treasure Island (FL): StatPearls Publishing.
  53. Murphy, C. *et al.* (2014) 'Management of Large Erupting Complex Odontoma in Maxilla', *Case Reports in Pediatrics*, pp. 1–3. doi: 10.1155/2014/963962.
  54. Muthu, K. (2017) 'Ossifying Fibroma of Maxilla', *Austin Journal of Dentistry*. doi: 10.26420/austinjdent.2017.1088.
  55. Nogueira, A. S. *et al.* (2014) 'Simultaneous occurrence of dentigerous cyst and residual cyst in the maxilla', *Brazilian journal of otorhinolaryngology*, 80(1), pp. 88–89.
  56. de Oliveira, M. A. *et al.* (2019) 'The Importance of Histopathological

Examination to the Final Diagnosis of Peripheral Odontogenic Tumors: A Case Report of a Peripheral Odontoma', *Case reports in dentistry*, 2019, p. 9712816.

57. Packiri, S., Gurunathan, D. and Selvarasu, K. (2017) 'Management of Paediatric Oral Ranula: A Systematic Review', *Journal of clinical and diagnostic research: JCDR*, 11(9), pp. ZE06–ZE09.
58. Patil, S. B. *et al.* (2017) 'Comparison of Extended Nasolabial Flap Versus Buccal Fat Pad Graft in the Surgical Management of Oral Submucous Fibrosis: A Prospective Pilot Study', *Journal of maxillofacial and oral surgery*, 16(3), pp. 312–321.
59. Pattnaik, B. *et al.* (2017) 'Osteomyelitis of maxilla: A rarity', *Annals of International medical and Dental Research*. doi: 10.21276/aimdr.2017.3.3.de2.
60. Patturaja, K. and Pradeep, D. (2016) 'Awareness of Basic Dental Procedure among General Population', *Research Journal of Pharmacy and Technology*, p. 1349. doi: 10.5958/0974-360x.2016.00258.4.
61. Rajendran, A. and Sivapathasundharam, B. (2014) *Shafer's Textbook of Oral Pathology*. Elsevier Health Sciences.
62. Rao, T. D. and Santhosh Kumar, M. P. (2018) 'Analgesic Efficacy of Paracetamol Vs Ketorolac after Dental Extractions', *Research Journal of Pharmacy and Technology*, p. 3375. doi: 10.5958/0974-360x.2018.00621.2.
63. Regezi, J. A., Sciubba, J. and Jordan, R. C. K. (2012) *Oral Pathology - E-Book: Clinical Pathologic Correlations*. Elsevier Health Sciences.
64. Sadaksharam, J. and Murugesan, M. (2019) 'Osteomyelitis of Maxilla: A Rare Finding from a Radiologist Point of View', *Contemporary clinical dentistry*, 10(2), pp. 394–396.
65. Seehra, J., Sloan, P. and Oliver, R. J. (2009) 'Paget's disease of bone and osteonecrosis', *Dental update*, 36(3), pp. 166–8, 171–2.
66. Seo, Y.-K. *et al.* (2019) 'Diagnostic considerations in central odontogenic fibroma of the maxilla: 2 case reports', *Imaging science in dentistry*, 49(3), pp. 229–234.
67. Serindere, G. *et al.* (2019) 'Comparison of Head and Neck Cancer Distribution in Turkish and Syrian Populations', *Iranian journal of public health*, 48(10), pp. 1810–1816.
68. Shanmugam, V. U. *et al.* (2018) 'Fibrous Dysplasia of Maxilla - Surgical Sculpting for Aesthetics', *Journal of Medical Science And clinical Research*. doi: 10.18535/jmscr/v6i8.111.
69. Shimmura, H. *et al.* (2020) 'Metastasis of Papillary Thyroid Carcinoma to the Maxillary Sinus: Case Report and Review of the Literature', *Case reports in otolaryngology*, 2020, p. 4056901.
70. Silva, L. P. *et al.* (2018) 'A retrospective multicentre study of cystic lesions and odontogenic tumours in older people', *Gerodontology*, 35(4), pp. 325–332.
71. Sriram, G. and Shetty, R. P. (2008) 'Odontogenic tumors: a study of 250 cases in an Indian teaching hospital', *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, pp. e14–e21. doi: 10.1016/j.tripleo.2008.02.021.
72. Su, C.-W. *et al.* (2019) 'Spontaneous

osteosarcoma transformation of fibrous dysplasia in maxilla', *Journal of dental sciences*, 14(2), pp. 219–222.

73. Sweta, V. R., Abhinav, R. P. and Ramesh, A. (2019) 'Role of Virtual Reality in Pain Perception of Patients Following the Administration of Local Anesthesia', *Annals of maxillofacial surgery*, 9(1), pp. 110–113.
74. Tamiolakis, P. *et al.* (2019) 'Demographic and Clinical Characteristics of 5294 Jaw Cysts: A Retrospective Study of 38 Years', *Head and neck pathology*, 13(4), pp. 587–596.
75. Thakur, G. *et al.* (2011) 'Juvenile ossifying fibroma of the maxilla', *Case Reports*, pp. bcr0220113896–bcr0220113896. doi: 10.1136/bcr.02.2011.3896.
76. Thompson, L. D. R. and Bishop, J. A. (2017) *Head and Neck Pathology E-Book: A Volume in the Series: Foundations in Diagnostic Pathology*. Elsevier Health Sciences.
77. Wanjari, P. V., Wanjari, S. and Bhowate, R. (2011) 'Erupting complex odontoma of maxilla: A case report', *Journal of Pierre Fauchard Academy (India Section)*, pp. 65–68. doi: 10.1016/s0970-2199(11)50004-9.
78. Ward, P. H., Alley, C. and Owen, R. (1969) 'MONOSTOTIC FIBROUS DYSPLASIA OF THE MAXILLA', *The Laryngoscope*, pp. 1295–1306. doi: 10.1288/00005537-196907000-00007.
79. Xia, R.-H. *et al.* (2020) 'Ameloblastoma with mucous cells: A clinicopathological, BRAF mutation, and MAML2 rearrangement study', *Oral diseases*, 26(4), pp. 805–814.
80. Zhuoying, C. and Fengguo, Y. (2019) 'Huge erupted complex odontoma in maxilla', *Oral and Maxillofacial Surgery Cases*, p. 100096. doi: 10.1016/j.omsc.2019.100096.