

# Patient Comfort Level among Various Age Groups during Orthodontic Treatment- A Cross-Sectional Survey

**Running title:** Patient comfort level during orthodontic treatment

**Priadarsini T,**

Saveetha dental college and hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai - 77  
Email: 151501017.sdc@saveetha.com

**Naveen Kumar\***

Senior lecturer, Department of Orthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-77  
Email: naveenkumarm.sdc@saveetha.com

**Dhanraj Ganapathy**

Professor and HOD, Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University Chennai-77. Email: dhanraj@saveetha.com

## **Article Info**

**Volume 83**

**Page Number: 2375 - 2384**

**Publication Issue:**

**July-August 2020**

## **Abstract**

Orthodontic patients experience pain and discomfort to a varying degree during the course of treatment. One of the most important goals of orthodontic dental care is helping patients in their attempts to reach an acceptable level of satisfaction with their oral cavity and dentition. Dentofacial problems have known definitive effects on patient satisfaction and comfort with their appliance because it affects esthetics, performance, and function. The aim of this study was to assess the comfort level among various age groups during the orthodontic treatment. A questionnaire comprising 13 questions was created in an online survey platform and the link was shared to the patients. The patient data was collected from the case records of patients visiting Saveetha dental College for treatment. Statistical analysis was done using IBM SPSS software version 20.0. 44% of the respondents feel that their appliance is not at all comfortable and about 32% of the people are embarrassed to wear the appliance in public. Chi-square test was done for the association of gender and embarrassment of wearing appliances in public and the p value 0.001\* was found to be statistically significant. This concludes that more male patients feel embarrassed to wear the appliance in public and patients of age group 15-20 years feel that the appliance is more tight, eliciting pain.

**Keywords:** Adolescents ; Appliance ; Comfort ; Orthodontics ; Pain.

## **Article History**

**Article Received:** 06 June 2020

**Revised:** 29 June 2020

**Accepted:** 14 July 2020

**Publication:** 25 July 2020

## **INTRODUCTION**

An increasing number of patients seek orthodontic treatment because malocclusion has a significant effect on their quality of life (Jones, 1984) . Orthodontic appliances represent foreign objects inserted in a physically and psychologically sensitive area of the body (Sivamurthy and Sundari, 2016; Vikram *et al.*, 2017). It's usually obvious to others that they are being worn and it's possible that susceptible individuals may be self-conscious about wearing them in public (Dinesh *et al.*, 2013; Felicita, 2018) . Children in particular are subjected to social ridicule from their peers (Sergl, Klages and Pempera, 1992; Kamisetty *et al.*, 2015).

Yet orthodontic patients are expected to remain compliant and cooperative throughout the treatment which may last for several years (Krishnan, Pandian and Kumar S, 2015; Samantha, Sundari and Chandrasekhar, 2017) .The patients' psychological response to orthodontic treatment and their ability to adapt to appliances are clearly a very significant concern (Herren *et al.*, 1966). The discomfort of orthodontic treatment has a negative effect on patient compliance (Rubika, Sumathi Felicita and Sivambiga, 2015; Viswanath *et al.*, 2015) .

Among the adverse effects of orthodontic treatment, speech difficulty is one of the major complications (Jain, Kumar and Manjula, 2014; Felicita, 2017b) . Apart from the original speech problems caused by malocclusion, orthodontic appliances can also lead to speech disturbances because they are a foreign body in the oral cavity (Felicita, Chandrasekar and ShanthaSundari, 2012; Felicita, 2017). Specifically, orthodontic appliances often fit against the palate and the surface of the teeth, which affects the movement of the tongue and the space of the oral cavity, resulting in the distortion of some specific sounds .

In addition, it has been reported that various types of orthodontic appliances such as labial and lingual fixed appliances , tongue thrusts ,palatal expanders and Hawley retainers may influence speech clarity (Ramesh

Kumar *et al.*, 2011; Pandian, Krishnan and Kumar, 2018).

While many studies have taken the standpoint of operators, patient's own perceptions of their treatment have been rarely sought. The aim of this study was to evaluate the experience of patients wearing fixed and removable orthodontic appliances to and evaluate their comfort level.

## **MATERIALS AND METHODS**

### **Study design and setting**

A total of 100 patients who were undergoing orthodontic treatment in Saveetha dental college were randomly selected. The patients details were collected from the patient records and were contacted through for the distribution of the questionnaire created in an online survey platform.

### **Ethical Approval**

The study was commenced after approval from the institutional review board (Ethical approval number : SDC/SIHEC/2020/DIASDATA/0619-0320).

### **Informed consent**

A written informed consent was obtained from all the study subjects.

### **Subjects and Procedures**

The questionnaire was randomly distributed to 100 patients who were undergoing Orthodontic treatment at Saveetha dental college. The following data retrieved from the dental records: Patient's name , age, gender and contact number . The questionnaire comprised 13 questions which used a four point Likert type scale where each item responded is scored indicating the degree of agreement with the statement - Not at all, Little , Much and Very much.

### Statistical analysis

The statistical analysis was done using SPSS software version 21.0 (SPSS Inc., Chicago, IL, USA). Descriptive statistics (percentage and mean) and Inferential statistics (Chi Square test) were done.

### RESULTS AND DISCUSSION

Of the total hundred participants, 40% were female and 60% were male. They were split into two age groups 15 to 20 years which comprised 53% and 21 to 26 years which comprised 47% of the population.

Figure 1 day depicts the pie chart on the type of appliance where 64% were using fixed appliances and 36% used removable Orthodontic appliances.

In figure 2 appliance comfort has been depicted. Around 44% feel that the appliance is not at all comfortable followed by 40% of the respondents mentioning that the appliance is a little comfortable and 10% responding appliance is much comfortable and 6% responding at the appliance is very much comfortable.

Figure 3 was on appliance tightness. 51% of them feel that appliance is a little tight followed by 20% responding it is much tight, 19% as not at all tight and 10% as very much tight.

Figure 4 represents the interference of appliance in speech - 55% of the respondents state that it is very much interfering with the speaking followed by 25% much interference in speaking 12% as a little interference in speaking and 8% has stated that appliance has not at all interfered in their speech.

Figure 5 represents the response to being embarrassed to wear the appliance in public. 32% are very much embarrassed to wear the appliance in public, followed by 29% a little embarrassed to wear it in public, 26% much embarrassed and 13% not at all embarrassed about wearing the appliance in public. Figure 6 is the association of age and type of appliance which has a p value of about 0.652. In figure 7 the association of age and appliance comfort has been depicted. The p value was 0.028\* (statistically significant). In figure 8, for

the association of age with appliance tightness, p value was 0.12 ( $p > 0.05$ ; statistically insignificant). Figure 9 depicts the association of gender and being embarrassed to wear their appliance in public has been demonstrated. The p value for this is about 0.001\* which is statistically significant. Also, males were more embarrassed to wear the appliance in public than females.

For efficient clinical management of orthodontic patients, it is suggested to predict their behaviour and compliance during the treatment (Lewis and Brown, 1973). The amount of initial pain and discomfort experienced by the patient predicts the acceptance of the appliance and treatment in general (Oliver and Knapman, 1985). Patient self-confidence is affected by speech impairment during social interactions when focused on face, eyes and mouth (Sergl, Klages and Zentner, 1998). The case in our study was about 32% of the people were embarrassed to wear the appliance in public and in 55% of the patients they applied and interfered in their speech. It is to be noted that for an adolescent patient whose motivation for orthodontic treatment is guided primarily by the perception of their own appearances, during treatment they feel even more being the centre of attention of peers and acquaintances (Nanda and Kierl, 1992).

Another predictor of the discomfort during orthodontic treatment revealed in the present work was the value of dental aesthetics (Woolasset *al.*, 1988). The more pleasing the appliance looks aesthetically, the lower was the complaint intensity of feeling of tightness (Sergl and Zentner, 2000). So dentists might try to incorporate more aesthetic measures in the kids and adolescents in their removable appliance so that the kids might feel good using them. It is the role of the dentist to make their patients as comfortable as possible and also deliver the appliance in the correct functioning way.

### CONCLUSION

Prior to the orthodontic treatment, patients should be forewarned about the practical difficulties that may be induced by orthodontic appliances. Advice regarding the adaptation time should also be given. Dentists should advise the patients to speak more slowly and to accept some speech distortions during the period of adaptation. Patients can be counseled about the changes to expect with relation to pain, discomfort, speech, chewing, and swallowing immediately after an appliance is placed and up to a week after wearing it. The patient's psychological response to orthodontic treatment and ability to adapt to the appliance might be significantly improved with anticipatory guidance. Providing such information to patients can be expected to expedite the adaptation period and improve cooperation.

#### AUTHOR CONTRIBUTIONS

We thank all the above-mentioned authors for their valuable support and contribution to this study.


#### CONFLICTS OF INTEREST

There are no conflicts of interest.


#### REFERENCES

1. Dinesh, S. P. S. *et al.* (2013) 'An indigenously designed apparatus for measuring orthodontic force', *Journal of clinical and diagnostic research: JCDR*, 7(11), pp. 2623–2626.
2. Felicita, A. S. (2017a) 'Orthodontic management of a dilacerated central incisor and partially impacted canine with unilateral extraction - A case report', *The Saudi dental journal*, 29(4), pp. 185–193.
3. Felicita, A. S. (2017b) 'Quantification of intrusive/retraction force and moment generated during en-masse retraction of maxillary anterior teeth using mini-implants: A conceptual approach', *Dental press journal of orthodontics*, 22(5), pp. 47–55.
4. Felicita, A. S. (2018) 'Orthodontic extrusion of Ellis Class VIII fracture of maxillary lateral incisor–The sling shot method', *The Saudi dental journal*. Elsevier. Available at: <https://www.sciencedirect.com/science/article/pii/S1013905218302116>.
5. Felicita, A. S., Chandrasekar, S. and Shanthasundari, K. K. (2012) 'Determination of craniofacial relation among the subethnic Indian population: a modified approach - (Sagittal relation)', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 23(3), pp. 305–312.
6. Herren, P. *et al.* (1966) '[Teacher's questionnaire--a way of finding out psychologic factors in orthodontic diagnosis]', *Fortschritte der Kieferorthopadie*, 27(2), pp. 160–176.
7. Jain, R. K., Kumar, S. P. and Manjula, W. S. (2014) 'Comparison of intrusion effects on maxillary incisors among mini implant anchorage, j-hook headgear and utility arch', *Journal of clinical and diagnostic research: JCDR*, 8(7), pp. ZC21–4.
8. Jones, M. L. (1984) 'An investigation into the initial discomfort caused by placement of an archwire', *European journal of orthodontics*, 6(1), pp. 48–54.
9. Kamisetty, S. K. *et al.* (2015) 'SBS vs Inhouse Recycling Methods-An Invitro Evaluation', *Journal of clinical and diagnostic research: JCDR*, 9(9), pp. ZC04–8.
10. Krishnan, S., Pandian, S. and Kumar S, A. (2015) 'Effect of bisphosphonates on orthodontic tooth movement-an update', *Journal of clinical and diagnostic research: JCDR*, 9(4), pp. ZE01–5.
11. Lewis, H. G. and Brown, W. A. (1973) 'The attitude of patients to the wearing of a removable orthodontic appliance', *British dental journal*, 134(3), pp. 87–90.
12. Nanda, R. S. and Kierl, M. J. (1992) 'Prediction of cooperation in orthodontic treatment', *American journal of orthodontics and dentofacial orthopedics: official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*, 102(1), pp. 15–21.


13. Oliver, R. G. and Knapman, Y. M. (1985) 'Attitudes to orthodontic treatment', *British journal of orthodontics*, 12(4), pp. 179–188.
14. Pandian, K. S., Krishnan, S. and Kumar, S. A. (2018) 'Angular photogrammetric analysis of the soft-tissue facial profile of Indian adults', *Indian journal of dental research: official publication of Indian Society for Dental Research*, 29(2), pp. 137–143.
15. Ramesh Kumar, K. R. *et al.* (2011) 'Depth of resin penetration into enamel with 3 types of enamel conditioning methods: A confocal microscopic study', *American journal of orthodontics and dentofacial orthopedics: official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*, 140(4), pp. 479–485.
16. Rubika, J., Sumathi Felicita, A. and Sivambiga, V. (2015) 'Gonial Angle as an Indicator for the Prediction of Growth Pattern', *World Journal of Dentistry*, pp. 161–163. doi: 10.5005/jp-journals-10015-1334.
17. Samantha, C., Sundari, S. and Chandrasekhar, S. (2017) 'Comparative evaluation of two Bis-GMA based orthodontic bonding adhesives-A randomized clinical trial', *Journal of clinical and*. ncbi.nlm.nih.gov. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5449915/>.
18. Serogl, H. G., Klages, U. and Pempera, J. (1992) 'On the prediction of dentist-evaluated patient compliance in orthodontics', *European journal of orthodontics*, 14(6), pp. 463–468.
19. Serogl, H. G., Klages, U. and Zentner, A. (1998) 'Pain and discomfort during orthodontic treatment: causative factors and effects on compliance', *American journal of orthodontics and dentofacial orthopedics: official publication of the American Association of Orthodontists, its constituent societies, and the American Board of Orthodontics*, 114(6), pp. 684–691.
20. Serogl, H. G. and Zentner, A. (2000) 'Predicting patient compliance in orthodontic treatment', *Seminars in orthodontics*, 6(4), pp. 231–236.
21. Sivamurthy, G. and Sundari, S. (2016) 'Stress distribution patterns at mini-implant site during retraction and intrusion—a three-dimensional finite element study', *Progress in orthodontics*, 17(1), p. 4.
22. Vikram, N. R. *et al.* (2017) 'Ball Headed Mini Implant', *Journal of clinical and diagnostic research: JCDR*, 11(1), pp. ZL02–ZL03.
23. Viswanath, A. *et al.* (2015) 'Obstructive sleep apnea: awakening the hidden truth', *Nigerian journal of clinical practice*, 18(1), pp. 1–7.
24. Woolass, K. F. *et al.* (1988) 'The prediction of patient co-operation in orthodontic treatment', *European journal of orthodontics*, 10(3), pp. 235–243.




**Figure 1:** Pie chart depicts the type of appliance used .36% of them used removable appliances (red) and 64% of them wore fixed appliances (blue).




**Figure 2:** Pie chart depicts the appliance comfort . 40% report the appliance is little comfortable (blue) , 10% report the appliance to be much comfortable (red) , 6%report the appliance to be very much comfortable (orange) and 44% reported the appliance to be not at all comfortable(green) .




**Figure 3:** Pie chart denotes the appliance tightness. Half of them ie, 51% reported the appliance to be a little tight (blue) , 20% reported the appliance to be much tight (red) , 10% reported the appliance to be very much tight (orange) and the remaining 19% reported the appliance to be not at all tight (green).




**Figure 4:** Pie chart denotes the appliance in speech interference. More than half ie, 55% reported that the appliance interferes in speech a little (blue) , 25% reported the appliance interference in speech to be much (red) , 12% reported the appliance interference in speech to be very much (orange) and the remaining 8% reported the appliance does not interfere with speech at all (green).




**Figure 5:** Pie chart depicts the response of embarrassment to wear the appliance in public. 29% of the respondents reported that they are a little embarrassed to wear the appliance in public (blue), 26% reported being much embarrassed (red) , 32% reported being very much embarrassed (orange) and the remaining 13% reported that they are not at all embarrassed to wear the appliance in public (green).




**Figure 6:** Bar graph depicts the association of age and type of appliance . X axis represents the age group and Y axis denotes the type of appliance. Chi-square test was done and was found to be statistically insignificant [Pearson’s Chi - Square value is 0.20;  $p=0.652$  ( $p>0.05$ )] More patients with fixed appliances fall into the age group of 15-20 years than 21-26 years , and those who use removable appliances are the same in number for both the age groups.



**Figure 7 :**Bar graph depicting the association of age and appliance comfort , X axis represents the age group and Y axis denotes the responses to appliance comfort. Chi-square test was done and was found to be statistically significant [Pearson’s Chi - Square value is 0.11;  $p=0.028^*$  ( $p<0.05$ )]. Patients in the age group of 15-20 years have responded the most for appliances to be not at all comfortable .



**Figure 8:** Bar graph depicting the association of age and appliance feeling tight. X axis denotes the age group and Y axis denotes the response for appliance tightness. Chi-square test was done and was found to be statistically insignificant [Pearson's Chi - Square value is 2.71;  $p = 0.12$  ( $p < 0.05$ )]. Patients in the age group of 21-26 years have responded the most for appliances to be very much tight.



**Figure 9:** Bar graph depicting the association of gender and response to embarrassment of wearing appliances in public. X axis denotes the gender and Y axis denotes the response to embarrassment to wear appliances in public. Chi-square test was done and was found to be statistically significant [Pearson's Chi - Square value is 0.26;  $p = 0.001^*$  ( $p < 0.05$ )]. Males are more embarrassed to wear the appliance in public than females.